



## Remarks about a fractional Choquard equation: ground state, regularity and polynomial decay

Pedro Belchior, Hamilton Bueno, Gilberto Assis Pereira,

Departamento de Matemática - UFMG

E-mail: pedrobelchior18@gmail.com, hamilton@mat.ufmg.br, gilbertoapereira@yahoo.com.br

Olímpio Miyagaki

Departamento de Matemática - UFJF

E-mail: ohmiyagaki@gmail.com

We consider a Choquard-type equation in a fractional  $(s, p)$ -Laplacian setting:

$$(-\Delta_p)^s u + A|u|^{p-2}u = \left( \frac{1}{|x|^\mu} * F(u) \right) f(u) \quad \text{in } \mathbb{R}^N, \quad (1)$$

where  $A$  is a positive constant,  $0 < \mu < N$ ,  $F(t) = \int_0^t f(r)dr$  and  $(-\Delta_p)^s$  denotes the  $(s, p)$ -Laplacian operator.

In (1), we assume that  $f$  is a  $C^1$  function that satisfies

$$(f1) \quad \lim_{t \rightarrow 0} \frac{|f(t)|}{t^{p-1}} = 0;$$

$$(f2) \quad \lim_{t \rightarrow \infty} \frac{f(t)}{t^{q-1}} = 0 \text{ for some } p < q < \frac{p^*}{2} \left( 2 - \frac{\mu}{N} \right), \text{ where}$$

$$p_s^* = \frac{Np}{N - sp} \text{ with } sp < N;$$

$$(f3) \quad f'(t)t^2 - (p-1)f(t)t > 0 \text{ for all } t > 0.$$

Since we are looking for positive solutions, we suppose that  $f(t) = 0$  for  $t \leq 0$ .

We summarize our results:

**Theorem 1.** *Suppose that, for  $p < q < (N - \mu)p/(N - p)$ ,  $0 < \mu < N$ , conditions (f1)-(f3) are valid. Then, for any  $A > 0$ , problem (1) has a non-negative ground state solution  $v$ .*

The first result was shown by proving that the natural energy functional attached to this problem satisfy the mountain pass geometry. By introducing the Nehari manifold, we show that the minimizing sequence converges strongly.

Supposing that  $0 < \mu < p$ , we adapt ideas from Brasco, Mosconi e Squassina [2, Proposition 3.2] and prove additionally that the ground state solution is continuous:

**Theorem 2.** *Let  $v \in W^{s,p}(\mathbb{R}^N)$  be a non-negative solution of problem (1). Then  $v \in L^\infty(\mathbb{R}^N) \cap C^0(\mathbb{R}^N)$ .*

Finally, we prove that the ground state solution has polynomial decay:

**Theorem 3.** *Suppose that (f1)-(f3) hold and  $0 < \mu < p < q < \frac{p^*}{2} (2 - \frac{\mu}{N})$ . Then, there exist constants  $\rho > 0$  and  $C > 0$  such that the solution obtained in Theorem 1 satisfies*

$$v(x) \leq \frac{C}{|x|^{\frac{N-sp}{p-1}}} \quad \forall x > \rho.$$

## Referências

- [1] C. O. Alves and M. Yang: *Investigating the multiplicity and concentration behaviour of solutions for a quasi-linear Choquard equation via the penalization method*, Proc. R. Soc. Edinb. **146A** (2016), 23-58.
- [2] L. Brasco, S. Mosconi and M. Squassina: *Optimal Decay of extremal for the fractional Sobolev Inequality*, Calc. Var. **55** (2016): 23. doi:10.1007/s00526-016-0958-y
- [3] L. Brasco and E. Parini: *The second eigenvalue of the fractional  $p$ -Laplacian*, Advances in Calculus of Variations **9** 4, 323-355, ISSN (Online) 1864-8266, ISSN (Print) 1864-8258, doi:10.1515/acv-2015-0007
- [4] F. Demengel and G. Demengel: *Functional Spaces for the Theory of Elliptic Partial Differential Equations*, Springer, London, 2012.
- [5] E. Di Nezza, G. Palatucci and E. Valdinoci: *Hitchhikers guide to the fractional Sobolev spaces*, Bull. Sci. Math. **136** (2012), no. 5, 521-573.
- [6] P. Felmer, A. Quaas and J. Tan: *Positive solutions of the nonlinear Schrödinger equation with the fractional Laplacian*, Proc. Roy. Soc. Edinburgh Sect. A **142** (2012), no. 6, 1237-1262.

**Agradecimentos:** *Agradecemos à CAPES, CNPq e FAPEMIG pelo apoio financeiro.*